

Variables	BIG 1	BIG 2	BIG 2 a	BIG 3
Neurologic Exam	Normal	Normal	GCS 13-14	Focal Neurologic Deficit, Abnormal Pupillary Exam, GCS ≤ 12
Intoxication	No	Yes / No	Yes / No	Yes / No
Anticoagulation / Antiplatelet	No	No Warfarin with INR < 1.4	Aspirin alone Dipyridimole Warfarin with INR 1.5-1.6, Prophylactic dose of anticoagulation *see Management for initial CT neg	Clopidogrel, Ticagrelor, Prasugrel, Warfarin with INR ≥ 1.7, any Novel anticoagulants, Therapeutic dose of anticoagulation *see Management for initial CT neg
Skull Fracture	No	Non-Displaced	Non -displaced	Displaced / Open
Subdural Hematoma (SDH) Axial Thickness	< 4mm	5-7 mm	5-7 mm	≥ 8 mm
Epidural Hematoma (EDH) Axial Thickness	< 4mm	5-7 mm	5-7 mm	≥ 8 mm
Intraparenchymal Hemorrhage (IPH) / Contusion	< 4mm 1 location	5-7 mm 2 locations	5-7 mm 2 locations	≥ 8 mm Multiple Locations
Subarachnoid Hemorrhage (SAH)	None or Trace	Localized over Convexity	Localized in 2 locations/ Convexity	Scattered/Diffuse/Basal Cisterns, or Aneurysmal Pattern
Intraventricular Hemorrhage (IVH)	No	No	Trace / Dependent Occipital Horns	Yes

THERAPEUTIC PLAN

Variables	BIG 1	BIG 2	BIG 2 a	BIG 3
Hospitalization	Consider observation for 6 hours vs. Home with responsible adult	23 Hour Observation	23 Hour Observation	Yes
Repeat Head CT	No	No	Reasonable but not mandatory	Yes
Neurosurgery Consult	Consider	Consider	Yes	Yes
Transfer to Level I Trauma Center	No	No	Not Necessary	Yes